

## PROBLEMS ON RECURRENCES

1. Let  $T_0 = 2, T_1 = 3, T_2 = 6$ , and for  $n \geq 3$ ,

$$T_n = (n + 4)T_{n-1} - 4nT_{n-2} + (4n - 8)T_{n-3}.$$

The first few terms are: 2, 3, 6, 14, 40, 152, 784, 5168, 40576, 363392. Find, with proof, a formula for  $T_n$  of the form  $T_n = A_n + B_n$ , where  $\{A_n\}$  and  $\{B_n\}$  are well-known sequences.

2. Define  $u_n$  by  $u_0 = 0, u_1 = 4$ , and  $u_{n+2} = \frac{6}{5}u_{n+1} - u_n$ . Show that  $|u_n| \leq 5$  for all  $n$ .
3. Prove or disprove that there exists a positive real number  $u$  such that  $\lfloor u^n \rfloor - n$  is an even integer for all positive integers  $n$ .
4. Show that the next integer above  $(\sqrt{3} + 1)^{2n}$  is divisible by  $2^{n+1}$ .
5. Let  $n$  be a positive integer and let  $a_1, \dots, a_{n-1}$  be arbitrary real numbers. Define the sequences  $u_0, \dots, u_n$  and  $v_0, \dots, v_n$  inductively by  $u_0 = u_1 = v_0 = v_1 = 1$ , and  $u_{k+1} = u_k + a_k u_{k-1}$ ,  $v_{k+1} = v_k + a_{n-k} v_{k-1}$  for  $k = 1, \dots, n-1$ .

Prove that  $u_n = v_n$ .

6. Let  $a_0, a_1, \dots$  be an arbitrary sequence of positive integers, and  $p_0 = 1, q_0 = 0, p_1 = a_0, q_1 = 1$ . Consider the recurrence

$$p_{n+2} = a_{n+1}p_{n+1} + p_n,$$

$$q_{n+2} = a_{n+1}q_{n+1} + q_n.$$

Show that  $p_n, q_n$  are coprime for any  $n \geq 0$ .

7. Let  $Q_0(x) = 1, Q_1(x) = x$ , and

$$Q_n(x) = \frac{(Q_{n-1}(x))^2 - 1}{Q_{n-2}(x)}$$

for all  $n \geq 2$ . Show that, whenever  $n$  is a positive integer,  $Q_n(x)$  is equal to a polynomial with integer coefficients.

8. For a positive integer  $n$  and any real number  $c$ , define  $x_k$  recursively by  $x_0 = 0, x_1 = 1$ , and for  $k \geq 0$ ,

$$x_{k+2} = \frac{cx_{k+1} - (n-k)x_k}{k+1}.$$

Fix  $n$  and then take  $c$  to be the largest value for which  $x_{n+1} = 0$ . Find  $x_k$  in terms of  $n$  and  $k, 1 \leq k \leq n$ .

9. Solve the recurrence

$$(n+1)(n+2)a_{n+2} - 3(n+1)a_{n+1} + 2a_n = 0,$$

with the initial conditions  $a_0 = 2, a_1 = 3$ .

10. Define  $u_0 = 1$  and for  $n \geq 0$ ,

$$2u_{n+1} = \sum_{k=0}^n \binom{n}{k} u_k u_{n-k}.$$

Find a simple expression for  $u_n$ .

11. Given some distinct positive integers  $a_1, \dots, a_n$ . Two players are playing a game where there are  $m$  stones on the table at the beginning, and each player takes turn to choose a number  $i$  from 1 to  $n$  and take  $a_i$  stones from the table. A player loses if there is no valid move, i.e. there are not  $\min a_i$  stones to take away from the table. Let  $f(m)$  be 1 if the first player has a winning strategy, and 0 if the second player has a winning strategy. Show that  $f$  is eventually periodic.
12. Let  $a_1 < a_2$  be two given integers. For any integer  $n \geq 3$ , let  $a_n$  be the smallest integer which is larger than  $a_{n-1}$  and can be uniquely represented as  $a_i + a_j$ , where  $1 \leq i < j \leq n - 1$ . Given that there are only a finite number of even numbers in  $\{a_n\}$ , prove that the sequence  $\{a_{n+1} - a_n\}$  is eventually periodic, i.e. that there exist positive integers  $T, N$  such that for all integers  $n > N$ , we have

$$a_{T+n+1} - a_{T+n} = a_{n+1} - a_n.$$

13. Let  $1, 2, 3, \dots, 2005, 2006, 2007, 2009, 2012, 2016, \dots$  be a sequence defined by  $x_k = k$  for  $k = 1, 2, \dots, 2006$  and  $x_{k+1} = x_k + x_{k-2005}$  for  $k \geq 2006$ . Show that the sequence has 2005 consecutive terms each divisible by 2006.
14. Let  $a_0, a_1, \dots$  be a sequence such that  $a_0 = 1, a_1 = 2$  and  $a_{n+2} = 4a_{n+1} - a_n$  for all  $n \geq 0$ . Show that  $a_m | a_{(2k+1)m}$  for all nonnegative integers  $m, k$ .
15. Let  $a_0 = 2, a_1 = -3$  and for all  $n \geq 2$ ,

$$a_n = \sum_{i=0}^{n-1} (2i - 1)a_{n-i-1}.$$

Find a closed form for  $a_n$  in terms of  $n$ .

16. Solve the recurrence:  $a_0 = 1, a_1 = 2019$  and

$$a_{n+2}a_n = a_{n+1}(a_n + a_{n+1})$$

for all  $n \geq 0$ .

17. Solve the first order recursion given by  $x_0 = 1$  and  $x_n = 1 + (1/x_{n-1})$ . Does  $\{x_n\}$  approach a limiting value as  $n$  increases?
18. Let  $a_0 = 5/2$  and  $a_k = a_{k-1}^2 - 2$  for  $k \geq 1$ . Compute

$$\prod_{i=0}^{\infty} \left(1 - \frac{1}{a_i}\right)$$

in closed form.

19. Given a real number  $a$ , we define a sequence by  $x_0 = 1, x_1 = x_2 = a$ , and  $x_{n+1} = 2x_n x_{n-1} - x_{n-2}$  for  $n \geq 2$ . Prove that if  $x_n = 0$  for some  $n$ , then the sequence is periodic.
20. Let  $k$  be an integer greater than 1. Suppose that  $a_0 > 0$ , and define

$$a_{n+1} = a_n + \frac{1}{\sqrt[k]{a_n}}$$

for  $n > 0$ . Evaluate

$$\lim_{n \rightarrow \infty} \frac{a_n^{k+1}}{n^k}.$$

21. Let  $a_0, a_1, \dots$  be any sequence of integers where  $a_0 = 0$  and

$$a_{n+2} = ca_{n+1} + da_n$$

for all  $n \geq 0$  where  $c, d$  are some integers. Show that for any prime  $p$  that is not a factor of  $d$ , there exists  $1 \leq i \leq p + 1$  such that  $p|a_i$ .

If  $c = 6$  and  $d = -1$ , show furthermore that for any odd prime  $p$  there exists  $1 \leq i < p$  such that  $p|a_i$ .

22. Let  $a_0, a_1, \dots$  be a sequence of integers where  $a_0 = 0$  and

$$a_{n+2} = ca_{n+1} + a_n$$

for all  $n \geq 0$  where  $c$  is a given integer. By pigeon hole principle, one can show that for any prime  $p$ , there exists  $T \leq p^2$  such that

$$a_{n+T} \equiv a_n \pmod{p}$$

holds for all  $n \geq 0$ . Show a much stronger result: for any prime  $p$  that is sufficiently large, there exists  $T \leq 2p + 2$  such that

$$a_{n+T} \equiv a_n \pmod{p} \quad \forall n \geq 0.$$