

## PROBLEMS ON GENERATING FUNCTIONS

1. Let  $a_n = (n^2 + 1)3^n$ . Compute  $y = \sum_{n \geq 0} a_n x^n$ .

2. Compute

$$y = x + \frac{2}{3}x^3 + \frac{24}{35}x^5 + \frac{246}{357}x^7 + \cdots.$$

3. Given  $a_0 = 2$ ,  $a_1 = 3$ , and

$$(n+1)(n+2)a_{n+2} - 3(n+1)a_{n+1} + 2a_n = 0,$$

for  $n \geq 0$ , compute  $y = \sum_{n \geq 0} a_n x^n$ .

4. Given  $a_0 = 1$  and  $a_{n+1} = (n+1)a_n - \binom{n}{2}a_{n-2}$  for  $n \geq 0$ , compute  $y = \sum_{n \geq 0} a_n \frac{x^n}{n!}$ .

5. Let  $k$  be a positive integer and let  $m = 6k - 1$ . Let

$$S(m) = \sum_{j=1}^{2k-1} (-1)^{j+1} \binom{m}{3j-1}.$$

For example with  $k = 3$ ,

$$S(17) = \binom{17}{2} - \binom{17}{5} + \binom{17}{8} - \binom{17}{11} + \binom{17}{14}.$$

Prove that  $S(m)$  is never zero.

6. For nonnegative integers  $n$  and  $k$ , define  $Q(n, k)$  to be the coefficient of  $x^k$  in the expansion of  $(1 + x + x^2 + x^3)^n$ . Prove that

$$Q(n, k) = \sum_{j=0}^n \binom{n}{j} \binom{n}{k-2j}.$$

7. Let  $a_{m,n}$  denote the coefficient of  $x^n$  in the expansion of  $(1 + x + x^2)^m$ . Prove that for all  $k \geq 0$ ,

$$0 \leq \sum_{i=0}^{\lfloor 2k/3 \rfloor} (-1)^i a_{k-i,i} \leq 1.$$

8. Consider the power series expansion

$$\frac{1}{1 - 2x - x^2} = \sum_{n=0}^{\infty} a_n x^n.$$

Prove that, for each integer  $n \geq 0$ , there is an integer  $m$  such that  $a_n^2 + a_{n+1}^2 = a_m$ .

9. Let  $A = \{(x, y) : 0 \leq x, y \leq 1\}$ . For  $(x, y) \in A$ , let

$$S(x, y) = \sum_{\frac{1}{2} \leq \frac{m}{n} \leq 2} x^m y^n,$$

where the sum ranges over all pairs  $(m, n)$  of positive integers satisfying the indicated inequalities. Evaluate

$$\lim_{(x,y) \rightarrow (1,1), (x,y) \in A} (1 - xy^2)(1 - x^2y)S(x, y).$$

10. For a set  $S$  of nonnegative integers, let  $r_S(n)$  denote the number of ordered pairs  $(s_1, s_2)$  such that  $s_1 \in S$ ,  $s_2 \in S$ ,  $s_1 \neq s_2$ , and  $s_1 + s_2 = n$ . Is it possible to partition the nonnegative integers into two sets  $A$  and  $B$  in such a way that  $r_A(n) = r_B(n)$  for all  $n$ ?
11. For positive integers  $m$  and  $n$ , let  $f(m, n)$  denote the number of  $n$ -tuples  $(x_1, x_2, \dots, x_n)$  of integers such that  $|x_1| + |x_2| + \dots + |x_n| \leq m$ . Show that  $f(m, n) = f(n, m)$ .
12. Let  $S_n$  denote the set of all permutations of the numbers  $1, 2, \dots, n$ . For  $\pi \in S_n$ , let  $\sigma(\pi) = 1$  if  $\pi$  is an even permutation and  $\sigma(\pi) = -1$  if  $\pi$  is an odd permutation. Also, let  $\nu(\pi)$  denote the number of fixed points of  $\pi$ . Find

$$\sum_{\pi \in S_n} \frac{\sigma(\pi)}{\nu(\pi) + 1}.$$

13. Let  $S$  be the set of sequences of length 2018 whose terms are in the set  $\{1, 2, 3, 4, 5, 6, 10\}$  and sum to 3860. Prove that the cardinality of  $S$  is at most

$$2^{3680} \left( \frac{2018}{2048} \right)^{2018}.$$

14. Suppose that  $\mathbb{Z}$  is written as a disjoint union of  $n < \infty$  arithmetic progressions, with differences  $d_1 \geq d_2 \geq \dots \geq d_n \geq 1$ . Show that  $d_1 = d_2$ .
15. Solve the following equation for the power series  $F(x, y) = \sum_{m, n \geq 0} a_{mn} x^m y^n$ , where  $a_{mn} \in \mathbb{R}$ :

$$(xy^2 + x - y)F(x, y) = xF(x, 0) - y.$$

The point is to make sure that your solution has a power series expansion at  $(0, 0)$ .

16. Find a simple description of the coefficients  $a_n \in \mathbb{Z}$  of the power series  $F(x) = x + a_2 x^2 + a_3 x^3 + \dots$  satisfying the functional equation

$$F(x) = (1 + x)F(x^2) + \frac{x}{1 - x^2}.$$

17. Consider the power series  $y = \sum_{n=0}^{\infty} \binom{3n}{n} x^n = 1 + 3x + 15x^2 + \dots$ . Show that

$$(27x - 4)y^3 + 3y + 1 = 0.$$

18. Find the unique power series  $y = 1 + \frac{1}{2}x + \frac{1}{12}x^2 - \frac{1}{720}x^4 + \frac{1}{30240}x^6 + \dots$  such that for all  $n \geq 0$ , the coefficient of  $x^n$  in  $y^{n+1}$  is equal to 1.
19. Find the unique power series  $y = 1 + x - \frac{1}{2}x^2 + \frac{2}{3}x^3 + \dots$  such that the constant term is 1, the coefficient of  $x$  is 1, and for all  $n \geq 2$  the coefficient of  $x^n$  in  $y^n$  is 0.
20. Let  $f(m, 0) = f(0, n) = 1$  and  $f(m, n) = f(m - 1, n) + f(m, n - 1) + f(m - 1, n - 1)$  for  $m, n > 0$ . Show that

$$\sum_{n=0}^{\infty} f(n, n) x^n = \frac{1}{\sqrt{1 - 6x + x^2}}.$$