

INEQUALITIES

1. For $p > 1$ and a_1, a_2, \dots, a_n positive, show that

$$\sum_{k=1}^n \left(\frac{a_1 + a_2 + \dots + a_k}{k} \right)^p < \left(\frac{p}{p-1} \right)^p \sum_{k=1}^n a_k^p.$$

2. If $a_n > 0$ for $n = 1, 2, \dots$, show that

$$\sum_{n=1}^{\infty} \sqrt[n]{a_1 a_2 \dots a_n} \leq e \sum_{n=1}^{\infty} a_n,$$

provided that $\sum_{n=1}^{\infty} a_n$ converges.

3. For $n = 1, 2, 3, \dots$ let

$$x_n = \frac{1000^n}{n!}.$$

Find the largest term of the sequence.

4. Suppose that a_1, a_2, \dots, a_n with $n \geq 2$ are real numbers greater than -1 , and all the numbers a_j have the same sign. Show that

$$(1 + a_1)(1 + a_2) \dots (1 + a_n) > 1 + a_1 + a_2 + \dots + a_n.$$

5. If a_1, \dots, a_{n+1} are positive real numbers with $a_1 = a_{n+1}$, show that

$$\sum_{i=1}^n \left(\frac{a_i}{a_{i+1}} \right)^n \geq \sum_{i=1}^n \frac{a_{i+1}}{a_i}.$$

6. Show that for any real numbers a_1, a_2, \dots, a_n ,

$$\left(\sum_{i=1}^n \frac{a_i}{i} \right)^2 \leq \sum_{i=1}^n \sum_{j=1}^n \frac{a_i a_j}{i+j-1}.$$

7. Let $y = f(x)$ be a continuous, strictly increasing function of x for $x \geq 0$, with $f(0) = 0$, and let f^{-1} denote the inverse function to f . If a and b are nonnegative constants, then show that

$$ab \leq \int_0^a f(x) dx + \int_0^b f^{-1}(y) dy.$$

8. Let a_1, a_2, \dots, a_n be real numbers. Show that

$$\min_{i < j} (a_i - a_j)^2 \leq M^2 (a_1^2 + \dots + a_n^2),$$

where

$$M^2 = \frac{12}{n(n^2 - 1)}.$$

9. Let f be a continuous function on the interval $[0, 1]$ such that $0 < m \leq f(x) \leq M$ for all x in $[0, 1]$. Show that

$$\left(\int_0^1 \frac{dx}{f(x)} \right) \left(\int_0^1 f(x) dx \right) \leq \frac{(m+M)^2}{4mM}.$$

10. Consider any sequence a_1, a_2, \dots of real numbers. Show that

$$\sum_{n=1}^{\infty} a_n \leq \frac{2}{\sqrt{3}} \sum_{n=1}^{\infty} \left(\frac{r_n}{n} \right)^{1/2}$$

where

$$r_n = \sum_{k=n}^{\infty} a_k^2.$$

(If the left-hand side of the inequality is ∞ , then so is the right-hand side.)

11. Show that

$$\frac{1}{(n-1)!} \int_n^{\infty} w(t) e^{-t} dt < \frac{1}{(e-1)^n},$$

where t is real, n is a positive integer, and

$$w(t) = (t-1)(t-2) \cdots (t-n+1).$$

12. Let $(a_n)_{n=1}^{\infty}$ and $(b_n)_{n=1}^{\infty}$ be two sequences of positive numbers. Show that the following statements are equivalent:

- There is a sequence $(c_n)_{n=1}^{\infty}$ of positive numbers such that $\sum_{n=1}^{\infty} \frac{a_n}{c_n}$ and $\sum_{n=1}^{\infty} \frac{c_n}{b_n}$ both converge.
- $\sum_{n=1}^{\infty} \sqrt{\frac{a_n}{b_n}}$

13. Suppose that a, b, c are real numbers in the interval $[-1, 1]$ such that $1 + 2abc \geq a^2 + b^2 + c^2$. Prove that $1 + 2(abc)^n \geq a^{2n} + b^{2n} + c^{2n}$ for all positive integers n .

14. Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ is a two times differentiable function satisfying $f(0) = 1, f'(0) = 0$ and for all $x \in [0, \infty)$, it satisfies

$$f''(x) - 5f'(x) + 6f(x) \geq 0$$

Prove that, for all $x \in [0, \infty)$,

$$f(x) \geq 3e^{2x} - 2e^{3x}$$

15. Let $f : [0, 1] \rightarrow \mathbb{R}$ be a continuous function satisfying $xf(y) + yf(x) \leq 1$ for every $x, y \in [0, 1]$.

- (a) Show that $\int_0^1 f(x) dx \leq \frac{\pi}{4}$.
 (b) Find such a function for which equality occurs.

16. For what pairs of positive real numbers (a, b) does the improper integral shown converge?

$$\int_b^{\infty} \left(\sqrt{\sqrt{x+a} - \sqrt{x}} - \sqrt{\sqrt{x} - \sqrt{x-b}} \right) dx$$

17. Let A be a positive real number. What are the possible values of $\sum_{j=0}^{\infty} x_j^2$, given that x_0, x_1, \dots are positive numbers for which $\sum_{j=0}^{\infty} x_j = A$?

18. Let $f(x)$ be a continuous real-valued function defined on the interval $[0, 1]$. Show that

$$\int_0^1 \int_0^1 |f(x) + f(y)| dx dy \geq \int_0^1 |f(x)| dx$$

19. For each continuous function $f : [0, 1] \rightarrow \mathbb{R}$, let $I(f) = \int_0^1 x^2 f(x) dx$ and $J(f) = \int_0^1 x (f(x))^2 dx$. Find the maximum value of $I(f) - J(f)$ over all such functions f .
20. Suppose that $f : [0, 1] \rightarrow \mathbb{R}$ has a continuous derivative and that $\int_0^1 f(x) dx = 0$. Prove that for every $\alpha \in (0, 1)$,

$$\left| \int_0^\alpha f(x) dx \right| \leq \frac{1}{8} \max_{0 \leq x \leq 1} |f'(x)|$$

21. For $m \geq 3$, a list of $\binom{m}{3}$ real numbers a_{ijk} ($1 \leq i < j < k \leq m$) is said to be area definite for \mathbb{R}^n if the inequality

$$\sum_{1 \leq i < j < k \leq m} a_{ijk} \cdot \text{Area}(\triangle A_i A_j A_k) \geq 0$$

holds for every choice of m points A_1, \dots, A_m in \mathbb{R}^n . For example, the list of four numbers $a_{123} = a_{124} = a_{134} = 1, a_{234} = -1$ is area definite for \mathbb{R}^2 . Prove that if a list of $\binom{m}{3}$ numbers is area definite for \mathbb{R}^2 , then it is area definite for \mathbb{R}^3 .

22. Let X_1, X_2, \dots be independent random variables with the same distribution, and let $S_n = X_1 + X_2 + \dots + X_n, n = 1, 2, \dots$. For what real numbers c is the following statement true:

$$\mathbb{P} \left(\left| \frac{S_{2n}}{2n} - c \right| \leq \left| \frac{S_n}{n} - c \right| \right) \geq \frac{1}{2}.$$

23. Let $H_k = \sum_{i=1}^k \frac{1}{i}$. Prove that the function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) = 1 + \sum_{n=1}^{\infty} \frac{x^n}{\prod_{k=1}^n H_k}$$

has no real zeros.

24. Let f be a continuous, nonnegative function on $[0, 1]$. Show that

$$\int_0^1 f(x)^3 dx \geq 4 \left(\int_0^1 x f(x)^2 dx \right) \left(\int_0^1 x^2 f(x) dx \right)$$

25. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be an infinitely differentiable function satisfying $f(0) = 0, f(1) = 1$, and $f(x) \geq 0$ for all $x \in \mathbb{R}$. Show that there exist a positive integer n and a real number x such that $f^{(n)}(x) < 0$.

26. Let $f = (f_1, f_2)$ be a function from \mathbb{R}^2 to \mathbb{R}^2 with continuous partial derivatives $\frac{\partial f_i}{\partial x_j}$ that are positive everywhere. Suppose that

$$\frac{\partial f_1}{\partial x_1} \frac{\partial f_2}{\partial x_2} - \frac{1}{4} \left(\frac{\partial f_1}{\partial x_2} + \frac{\partial f_2}{\partial x_1} \right)^2 > 0$$

everywhere. Prove that f is one-to-one.

27. Determine the greatest possible value of $\sum_{i=1}^{10} \cos(3x_i)$ for real numbers x_1, x_2, \dots, x_{10} satisfying $\sum_{i=1}^{10} \cos(x_i) = 0$.